

Thomas Mathew<sup>†</sup>

Joint work with: S.Zhai<sup>†</sup>, M. Flouri<sup>†</sup>, I.Bebu\*, B.Egan\*\*

<sup>†</sup>Department of Mathematics and Statistics, University of Maryland, Baltimore County

\*Biostatistics Center, George Washington University \*\*Department of Preventive Medicine and Biometrics, Uniformed Services University of the Health Sciences

## INTRODUCTION AND MOTIVATION

### MOTIVATION

Cost-effectiveness analysis is an integral part of health technology assessment, and addresses the question of whether a new treatment or other health care intervention offers good value for the money.

Cost is typically expressed in monetary terms, and the effectiveness is measured using variables such as survival time, Disability-Adjusted Life Years (DALYs), Quality-Adjusted Life Years (QALYs), et.

While evaluating treatments regarding their costs and health benefits, traditionally only population averages are compared. We wish to perform cost-effectiveness analysis, as it applies to the entire population. We propose to develop criteria that can bring out features not captured by the usual summary measures based on means; in particular, criteria involving medians, percentiles etc. This will enable us to answer questions such as: for a significant percentage of the population, is the increase in cost too large compared to the effectiveness gain? Statistical tolerance limits will be used to investigate such issues.

## BACKGROUND

- Consider the scenario where two treatments are compared using two groups of patients. Let  $C_i^j$  ( $E_i^j$ ) denote the cost (effect) of subject  $i$  in the  $j^{th}$  group,  $i = 1, 2, \dots, n_j, j = 1, 2$

- Traditional cost-effectiveness analysis is based on **Incremental Cost-Effectiveness Ratio (ICER)** and **Incremental Net Benefit (INB)**

$$ICER = \frac{\mu_C^1 - \mu_C^2}{\mu_E^1 - \mu_E^2}, \quad INB = \lambda(\mu_E^1 - \mu_E^2) - (\mu_C^1 - \mu_C^2)$$

$\mu_C^j$  ( $\mu_E^j$ ): mean of cost (effect) in the  $j^{th}$  group  
 $\lambda$ : Willingness-to-Pay amount

- ICER and INB are population parameters. The corresponding random variables are:

$$Y_{ICER} = \frac{C^1 - C^2}{E^1 - E^2}, \quad Y_{INB} = \lambda(E^1 - E^2) - (C^1 - C^2)$$

## SETUP

### Bivariate Normal

- Typically costs follow a log-normal distribution. However, the normal distribution is also used in practice.

- Under normality,  $\begin{pmatrix} C_i^j \\ E_i^j \end{pmatrix} \sim N\left(\begin{pmatrix} \mu_C^j \\ \mu_E^j \end{pmatrix}, \Sigma^j\right)$

- When the costs are log-normal, we assume  $\begin{pmatrix} \ln(C_i^j) \\ E_i^j \end{pmatrix} \sim N\left(\begin{pmatrix} \mu_C^j \\ \mu_E^j \end{pmatrix}, \Sigma^j\right)$


## MEDIAN AND THE ICER

The median of  $Y_{ICER}$  could be drastically different from the value of ICER. In other words, the traditional ICER could be unrepresentative of the population. In order to see this, consider the Lognormal-normal case with the following parameter values:

$$\mu^1 = \begin{pmatrix} \mu_C^1 \\ \mu_E^1 \end{pmatrix} = \begin{pmatrix} 8 \\ 10 \end{pmatrix}, \quad \Sigma^1 = \begin{pmatrix} 1.6 & -1.789 \\ -1.789 & 8 \end{pmatrix}$$

$$\mu^2 = \begin{pmatrix} \mu_C^2 \\ \mu_E^2 \end{pmatrix} = \begin{pmatrix} 6 \\ 8 \end{pmatrix}, \quad \Sigma^2 = \begin{pmatrix} 1.2 & -1.342 \\ -1.342 & 6 \end{pmatrix}$$

The following histogram shows the distribution of  $Y_{ICER}$ , along with the median and the true value of ICER. Clearly, the ICER is unrepresentative of the distribution.



## ONE-SIDED TOLERANCE LIMIT

- Tolerance limits are computed using random sample
- Lower tolerance limit,  $L$ , for  $Y_{INB}$ :  $P_{data}[P(Y_{INB} \geq L | \text{data}) \geq p] = 1 - \alpha$
- Upper tolerance limit,  $U$ , for  $Y_{ICER}$ :  $P_{data}[P(Y_{ICER} \leq U | \text{data}) \geq p] = 1 - \alpha$

We consider non-parametric tolerance limits based on order statistics [1]

- Let  $X = (X_1, \dots, X_N)$  be a sample from a distribution and  $(X_{(1)}, \dots, X_{(N)})$  the corresponding order statistics

- Want  $m$  such that  $P_{X_{(m)}}[P_X(X \geq X_{(m)}) \geq p] = 1 - \alpha$
- $X_{(m)}$  is the intended lower tolerance
- $m$  is defined as the largest integer such that

$$P(W \geq N - m + 1 | N, 1 - p) \geq 1 - \alpha$$

where  $W \sim Bin(N, 1 - p)$

- Similarly an upper tolerance limit can be defined

[1] Krishnamoorthy, K. and Mathew, T. (2009). Statistical Tolerance Regions: Theory, Applications, and Computation. Wiley, J. and Sons.


## PARAMETRIC BOOTSTRAP WITH CALIBRATION

Samples are not directly available from the distributions of  $Y_{ICER}$  and  $Y_{INB}$ , so we proceed as follows:

- Generate pairs  $\begin{pmatrix} C_i^{1*} \\ E_i^{1*} \end{pmatrix}, \begin{pmatrix} C_i^{2*} \\ E_i^{2*} \end{pmatrix}$  by parametric bootstrap,  $i = 1, 2, \dots, B_1$
- Compute  $Y_{ICER,i}^*, Y_{INB,i}^*, i = 1, 2, \dots, B_1$
- Use the order statistics based on  $Y_{ICER,i}^*$  and  $Y_{INB,i}^*$  to obtain tolerance limits

Coverage probabilities of the resulting tolerance limits are not satisfactory

- Use bootstrap calibration to improve the coverage
- Use  $B_2$  second level bootstrap samples



## SIMULATIONS

### Bivariate Normal

$$\mu^1 = \begin{pmatrix} \mu_{11} \\ 10 \end{pmatrix}, \quad \Sigma^1 = \begin{pmatrix} 500 & \sigma_{12}^{(1)} \\ \sigma_{21}^{(1)} & 8 \end{pmatrix}, \quad \mu^2 = \begin{pmatrix} 485 \\ 10 \end{pmatrix}, \quad \Sigma^2 = \begin{pmatrix} 450 & \sigma_{12}^{(2)} \\ \sigma_{21}^{(2)} & 6 \end{pmatrix}$$

where  $\sigma_{12}^{(1)} = \rho\sqrt{500 \times 8}$ ,  $\sigma_{12}^{(2)} = \rho\sqrt{450 \times 6}$ ,  $\lambda = 25$

One-sided lower tolerance limit for INB:

$n_1 = n_2 = 50, 150$ ,  $\mu_{11} = 500, 525$  and  $\rho = -0.5, 0.1$

### Lognormal-Normal

$$\mu^1 = \begin{pmatrix} \mu_{11} \\ 10 \end{pmatrix}, \quad \Sigma^1 = \begin{pmatrix} 1.6 & \sigma_{12}^{(1)} \\ \sigma_{21}^{(1)} & 8 \end{pmatrix}, \quad \mu^2 = \begin{pmatrix} 6 \\ 8 \end{pmatrix}, \quad \Sigma^2 = \begin{pmatrix} 1.2 & \sigma_{12}^{(2)} \\ \sigma_{21}^{(2)} & 6 \end{pmatrix}$$

where  $\sigma_{12}^{(1)} = \rho\sqrt{1.6 \times 8}$ ,  $\sigma_{12}^{(2)} = \rho\sqrt{1.2 \times 6}$ ,  $\lambda = 5000$

One-sided lower tolerance limit for INB:

$n_1 = n_2 = 50, 150$ ,  $\mu_{11} = 8, 10$  and  $\rho = -0.5, 0.1$

First level bootstrap sample size  $B_1 = 5000$ , second level bootstrap sample size  $B_2 = 1000$ , number of simulation runs  $B_0 = 1000$ .

## RESULTS

Bivariate Normal with $p = 0.9$ and $1 - \alpha = 0.95$					Lognormal-Normal with $p = 0.9$ and $1 - \alpha = 0.95$				
n	$\mu_{11}$	$\rho$	INB coverage	ICER coverage	n	$\mu_{11}$	$\rho$	INB coverage	ICER coverage
50	500	-0.5	0.9585	0.9545	50	6	-0.5	0.9450	0.9440
50	525	-0.5	0.9470	0.9630	50	8	-0.5	0.9430	0.9450
150	500	-0.5	0.9535	0.9595	150	6	-0.5	0.9445	0.9505
150	525	-0.5	0.9480	0.9575	150	8	-0.5	0.9515	0.9500
50	500	0.1	0.9505	0.9505	50	6	0.1	0.9525	0.9465
50	525	0.1	0.9455	0.9560	50	8	0.1	0.9475	0.9370
150	500	0.1	0.9500	0.9595	150	6	0.1	0.9485	0.9440
150	525	0.1	0.9560	0.9555	150	8	0.1	0.9475	0.9390

## EXAMPLE

### Canadian Implant Defibrillator Study (CIDS)<sup>[2]</sup>

Implant Cardioverter Defibrillator (ICD) vs. Amiodarone (Amd) in reducing the risk of death in survivors of Ventricular Tachycardia (VT).

$n_{ICD} = 212$  subjects,  $n_{Amd} = 218$  subjects,  $\lambda = 100000$

Bivariate Normal Data with:

$$\begin{pmatrix} \mu_C^1 \\ \mu_E^1 \end{pmatrix} = \begin{pmatrix} 87044 \\ 4.832 \end{pmatrix}, \quad \Sigma^1 = \begin{pmatrix} 8462152 & 125.8 \\ 125.8 & 0.02418 \end{pmatrix}, \quad \begin{pmatrix} \mu_C^2 \\ \mu_E^2 \end{pmatrix} = \begin{pmatrix} 38819 \\ 4.682 \end{pmatrix}, \quad \Sigma^2 = \begin{pmatrix} 6497962 & 20.42 \\ 20.42 & 0.0244 \end{pmatrix}$$

Point Estimates:

$$\hat{\Delta}_c = 48225, \quad \hat{\Delta}_e = 0.15, \quad \widehat{INB} = \lambda \hat{\Delta}_e - \hat{\Delta}_c, \quad \widehat{ICER} = \frac{\hat{\Delta}_c}{\hat{\Delta}_e}$$

### 100(1 - 2 $\alpha$ )% Confidence Interval

Using Fieller's theorem

$$\widehat{ICER} \left( 1 - z_{1-\alpha}^2 c \pm z_{1-\alpha} \sqrt{a + b - 2c - z_{1-\alpha}^2 (ab - c^2)} \right) / (1 - z_{1-\alpha}^2 a), \quad \widehat{INB} \pm z_{1-\alpha} \sqrt{u\lambda}$$

where

$$a = \frac{\hat{V}(\hat{\Delta}_e)}{\hat{\Delta}_e^2}, \quad b = \frac{\hat{V}(\hat{\Delta}_c)}{\hat{\Delta}_c^2}, \quad c = \frac{\hat{C}(\hat{\Delta}_e, \hat{\Delta}_c)}{\hat{\Delta}_e \hat{\Delta}_c}, \quad u\lambda = \lambda^2 \hat{V}(\hat{\Delta}_e) + \hat{V}(\hat{\Delta}_c) - 2\lambda \hat{C}(\hat{\Delta}_e, \hat{\Delta}_c)$$

- 95% Lower Confidence Limit for INB: -75785.45
- 95% Upper Confidence Limit for ICER: 83737.26

### Tolerance Limits

- $p = 0.9, 1 - \alpha = 0.95$  Lower Tolerance limit for  $Y_{INB}$ : -63846.16
- $p = 0.9, 1 - \alpha = 0.95$  Upper Tolerance limit for  $Y_{ICER}$ : 898636.2
- $p = 0.03, 1 - \alpha = 0.95$  Lower Tolerance limit for  $Y_{INB}$ : 103.9
- $p = 0.2, 1 - \alpha = 0.95$  Upper Tolerance limit for  $Y_{ICER}$ : -127004.8

## Conclusions

- 90% or more of individual INB values fall above: -63846.16
- 90% or more of individual ICER values fall below 898636.2
- Even though the average INB has a negative lower limit, at least 3% of the values are positive
- Even though the average ICER has a positive upper limit, at least 20% of the values are negative
- This shows that the traditional ICER analysis is inadequate for cost-effectiveness analysis

[2] Willan, A.R. and Briggs, A.H.(2006). Statistical analysis of cost-effectiveness data. Wiley:Chichester, UK.

## ACKNOWLEDGMENT

The hardware used in the computational studies is part of the UMBC High Performance Computing Facility (HPCF). The facility is supported by the U.S. National Science Foundation through the MRI program (grant nos. CNS-0821258 and CNS-1228778) and the SCREMS program (grant no. DMS-0821311), with additional substantial support from the University of Maryland, Baltimore County (UMBC). See [www.umbc.edu/hpcf](http://www.umbc.edu/hpcf) for more information on HPCF and the projects using its resources.